

2985. CHROMATIN DEFECTS INDUCED UPON KNOCKOUT OF MAJOR CHROMATIN REMODELING FACTOR ISWI ATPASE SMARCA5 IN MOUSE ARE SENSED VIA P53 PATHWAY AND BLOCK CELL CYCLE PROGRESSION

Zikmund T., Kokavec J., Savvulidi F., Turkova T., Paszekova H., Skoultchi A. I., Stopka T. (*Biocev, 1st Faculty of Medicine, Charles University, Prague – CZ; Department of Pathological Physiology, 1st Faculty of Medicine, Charles University in Prague – CZ; Department of Cell Biology, Albert Einstein College of Medicine, Bronx, NY – USA*)

Chromatin structure is a major prerequisite of proper developmental processes. Chromatin structure is established, maintained and disrupted by SWI/SNF2

superfamily of ATP-dependent helicases. Sensing of irregularities of chromatin structure is not fully understood. Subfamily of ISWI ATPases has been proposed to regulate developmental processes although the particular mechanisms of its activity were elusive. We herein utilized conditional deletion of major ISWI member, Smarca5, in murine early thymic progenitors. Phenotype of Smarca5 mutants is not indicative of being mediated via defective T-cell receptor (TCR) rearrangement or dysregulation of gene repair. Smarca5 deficiency resulted in proliferation defect together with activation of DNA damage pathway. Production of double Smarca5^{-/-} Trp53^{-/-} mutants partially restored proliferation blockade but neither rescued developmental disruption of the thymocyte development nor aggravated lymphomagenesis in the Trp53^{-/-} mutants.

SYMPOSIUM ON ADVANCES IN MOLECULAR HEMATOLOGY 3: STATE OF THE ART INVITED LECTURE

3003. DNA DAMAGE RESPONSE, AGING AND CANCER: MECHANISMS AND OPPORTUNITIES FOR TREATMENT

Bartek J. (*Danish Cancer Society Research Center, Copenhagen – DK; Institute of Molecular and Translational Medicine, Palacky University, Olomouc – CZ*)

Abstrakt sdělení není k dispozici.

KONFERENCE OŠETŘOVATELSTVÍ 2

2983. VÝBĚR OPTIMÁLNÍHO ŽILNÍHO VSTUPU S OHLEDEM NA SPECIFIKA HEMATOLOGICKÉHO A HEMATOONKOLOGICKÉHO PACIENTA

Maňásek V., Ďuraš J., Jelínek T. (*Komplexní onkologické centrum Nový Jičín, Společnost pro porty a permanentní katetry, Nový Jičín; Klinika hematoonkologie, Fakultní nemocnice, Ostrava*)

Hematologie a hematoonkologie jsou obory, které využívají dominantně parenterální formu léčby, a proto je správný výběr žilního přístupu stěžejní pro opti-

mální aplikaci léčiv. Využíváme přitom vstupů krátkodobých, střednědobých i dlouhodobých. Volba žilního přístupu je výsledkem rozvahy, hodnotící charakter a délku podávané léčby, vlastnosti pacienta a stav žilního systému. V případě indikace centrálního přístupu nejčastěji volíme mezi klasickou centrální kanylou (netunelizovanou), tunelizovanou centrální kanylou, PICC (centrální kanylou zaváděnou z periferie) a portem. Každý z těchto druhů vstupů má své výhody i nevýhody, a to nejen při inzerci, ale především při dlouhodobém používání, kdy je pacient obecně vystaven vyššímu riziku komplikací, a to především infekčních a trombo-